	Age (years)	Sex	Dermal invasion	Lymph node metastasis		- Diabetes mellitus	GLUT1 staining score			
Patient no.					Disease stage		Intraepidermal lesion	Dermal- invasive lesion	Metastatic lesion	
1	71	F	+	_	T2N0M0, Stage II	-	2	2	-	
2	69	М	+	-	T2N0M0, Stage II	-	0	4	-	
3	70	М	+	+	T2N1M1, Stage IV	-	1	4	4	
4	93	F	+	-	T2N0M0, Stage II	-	3	3	-	
5	76	F	+	-	T1N0M0, Stage I	-	2	2	-	
6	66	F	+	+	T1N2M1, Stage IV	-	2	3	4	
7	79	F	+	$+^{a}$	T2N1M1, Stage IV	-	0	1	NA	
8	70	F	+	$+^{a}$	T2N1M0, Stage IIIa	-	1	4	NA	
9	78	М	+	+	T1N1M0, Stage IIIa	-	1	2	2	
10	51	F	+	+	T1N2M0, Stage IIIb	NA	0	0	4	
11	76	М	+	+	T2N1M1, Stage IV	+	1	2	2	
12	63	F	NA	+	T1N2M1, Stage IV	NA	0	0	4	
12	82	F	+	+	T1N2M0, Stage IIIb	11A	0	1	4	
13	82 67	г М	+	+	T1N2M0, Stage III0	-	U	3	3 4	

Table SI. Clinicopathological parameters of 51 patients with extramammary Paget's disease.

15	75	F	+	-	T1N0M0, Stage I	-	1	1	-
16	72	F	+	-	T1N0M0, Stage I	-	2	3	-
17	69	F	+	-	T1N0M0, Stage I	+	1	2	-
18	87	М	+	-	T2N0M0, Stage II	-	0	2	-
19	83	М	+	-	T2N0M0, Stage II	-	2	3	-
20	74	М	+	+	T1N2M0, Stage IIIb	-	0	1	4
21	78	М	+	-	T1N0M0, Stage I	+	0	4	-
22	70	М	+	-	T1N0M0, Stage I	-	1	1	-
23	83	М	+	-	T1N0M0, Stage I	-	3	2	-
24	75	М	+	-	T1N0M0, Stage I	-	1	3	-
25	70	М	+	-	T1N0M0, Stage I	+	2	3	-
26	66	М	+	$+^{a}$	T1N2M1, Stage IV	-	1	4	NA
27	77	F	+	$+^{a}$	T1N1M0, Stage IIIa	+	2	2	NA
28	81	F	+	-	T1N0M0, Stage I	-	1	2	-
29	65	F	-	-	TisN0M0, Stage 0	-	1	-	-
30	72	М	-	-	TisN0M0, Stage 0	-	0	-	-
31	77	М	-	-	TisN0M0, Stage 0	+	1	-	-
32	73	F	-	-	TisN0M0, Stage 0	-	1	-	-
33	73	F	-	-	TisN0M0, Stage 0	-	0	-	-
34	70	М	-	-	TisN0M0, Stage 0	-	2	-	-
35	73	М	-	-	TisN0M0, Stage 0	-	1	-	-

36	82	F	-	-	TisN0M0, Stage 0	-	1	-	-
37	77	F	-	-	TisN0M0, Stage 0	+	1	-	-
38	59	F	-	-	TisN0M0, Stage 0	-	1	-	-
39	76	М	-	-	TisN0M0, Stage 0	+	0	-	-
40	77	F	-	-	TisN0M0, Stage 0	-	2	-	-
41	85	F	-	-	TisN0M0, Stage 0	+	0	-	-
42	71	М	-	-	TisN0M0, Stage 0	-	0	-	-
43	80	М	-	-	TisN0M0, Stage 0	-	0	-	-
44	72	F	-	-	TisN0M0, Stage 0	-	1	-	-
45	82	F	-	-	TisN0M0, Stage 0	-	0	-	-
46	65	М	-	-	TisN0M0, Stage 0	-	0	-	-
47	72	F	-	-	TisN0M0, Stage 0	-	1	-	-
48	80	М	-	-	TisN0M0, Stage 0	-	1	-	-
49	78	F	-	-	TisN0M0, Stage 0	-	0	-	-
50	69	М	-	-	TisN0M0, Stage 0	+	1	-	-
51	67	F	-	-	TisN0M0, Stage 0	-	0	-	-

<sup>a</sup>Histology not available. Disease stages were classified according to the proposed staging system by Ohara et al (1). F, female; M, male; GLUT1, glucose transporter isoform 1; NA, data not available.

## Reference

1. Ohara K, Fujisawa Y, Yoshino K, Kiyohara Y, Kadono T, Murata Y, Uhara H, Hatta N, Uchi H, Matsushita S, *et al.*: A proposal for a TNM staging system for extramammary Paget disease: Retrospective analysis of 301 patients with invasive primary tumors. J Dermatol Sci 83: 234-239, 2016.